## TABLE OF CONTENTS

1 Table of Contents
1.1 List of Tables
1.2 List of Figures
2 Introduction
2.1 Catalyst
2.2 Related Reports
2.3 Upcoming Reports
3 Epidemiology
3.1 Disease Overview
3.2 Risk Factors and Comorbidities
3.2.1 Family history is a strong predictor of obesity
3.2.2 Physical inactivity is an independent predictor of obesity
3.2.3 Excessive caloric intake doubles the risk for obesity
3.2.4 Hypertension is as high as 42% in obese adults
3.2.5 Dyslipidemia and type 2 diabetes are common comorbidities in obese patients
3.3 Global Trends
3.3.1 US
3.3.2 5EU
3.3.3 Japan
3.3.4 Brazil
3.3.5 Canada

- 3.4 Forecast Methodology
- 3.4.1 Sources Used
- 3.4.2 Forecast Assumptions and Methods: Prevalent Cases of Overweight, Obesity, Obesity by Class, and Comorbidities
- 3.4.3 Sources Not Used
- 3.5 Epidemiology Forecast of Overweight (2012–2022)
- 3.5.1 Prevalent Cases of Overweight
- 3.5.2 Age-Specific Prevalent Cases of Overweight
- 3.5.3 Sex-Specific Prevalent Cases of Overweight
- 3.5.4 Age-Standardized Prevalence of Overweight
- 3.6 Epidemiology Forecast of Obesity (2012–2022)
- 3.6.1 Prevalent Cases of Obesity
- 3.6.2 Age-Specific Prevalent Cases of Obesity
- 3.6.3 Sex-Specific Prevalent Cases of Obesity
- 3.6.4 Age-Standardized Prevalence of Obesity
- 3.7 Epidemiology Forecast of Obesity Class I (2012–2022)
- 3.7.1 Prevalent Cases of Obesity Class I
- 3.7.2 Age-Specific Prevalent Cases of Obesity Class I
- 3.7.3 Sex-Specific Prevalent Cases of Obesity Class I
- 3.7.4 Age-Standardized Prevalence of Obesity Class I
- 3.8 Epidemiology Forecast of Obesity Class II (2012–2022)
- 3.8.1 Prevalent Cases of Obesity Class II
- 3.8.2 Age-Specific Prevalent Cases of Obesity Class II
- 3.8.3 Sex-Specific Prevalent Cases of Obesity Class II

- 3.8.4 Age-Standardized Prevalence of Obesity Class II
- 3.9 Epidemiology Forecast of Obesity Class III (2012–2022)
- 3.9.1 Prevalent Cases of Obesity Class III
- 3.9.2 Age-Specific Prevalent Cases of Obesity Class III
- 3.9.3 Sex-Specific Prevalent Cases of Obesity Class III
- 3.9.4 Age-Standardized Prevalence of Obesity Class III
- 3.10 Epidemiology Forecast of Comorbidities among Adults with Overweight/Obesity (2012 and 2022)
- 3.10.1 Prevalent Cases of Diagnosed Diabetes among Adults with Overweight/Obesity
- 3.10.2 Prevalent Cases of Diagnosed Hypertension in Adults with Overweight/Obesity
- 3.10.3 Prevalent Cases of Dyslipidemia among Adults with Overweight/Obesity
- 3.11 Discussion
- 3.11.1 Conclusions on Epidemiological Trends
- 3.11.2 Limitations of the Analysis
- 3.11.3 Strengths of the Analysis
- 4 Appendix
- 4.1 Bibliography
- 4.2 About the Authors
- 4.2.1 Epidemiologists
- 4.2.2 Reviewers
- 4.2.3 Global Director of Epidemiology and Health Policy
- 4.2.4 Global Head of Healthcare
- 4.3 About GlobalData
- 4.4 About EpiCast

- 4.5 Contact Us
- 4.6 Disclaimer

## LIST OF TABLES

- Table 1: The WHO Classification System of Adult Overweight and Obesity According to BMI
- Table 2: Risk Factors and Comorbidities of Obesity
- Table 3: 9MM, Age-Adjusted and Crude Prevalence (%) of Obesity, by Sex, Ages =20 Years, 2008
- Table 4: JASSO and WHO Classifications of Obesity
- Table 5: Sources of Epidemiological Data Used for Forecasting the Prevalent Cases of Overweight
- Table 6: Sources of Epidemiological Data Used for Forecasting the Prevalent Cases of Obesity
- Table 7: Sources of Epidemiological Data Used for Forecasting the Prevalent Cases of Obesity Class I, Class II, and Class III
- Table 8: Sources of Epidemiological Data Used for Forecasting the Prevalent Cases of Comorbidities in Overweight/Obese
- Table 9: 9MM, Prevalent Cases of Overweight, Ages =18 Years, Both Sexes, N (Millions), 2012–2022
- Table 10: 9MM, Prevalent Cases of Overweight, By Age, Both Sexes, N (Millions), Row (%), 2012
- Table 11: 9MM, Prevalent Cases of Overweight, Ages = 18 Years, By Sex, N (Millions), Row (%), 2012
- Table 12: 9MM, Prevalent Cases of Obesity, Ages =18 Years, Both Sexes, N (Millions), 2012–2022
- Table 13: 9MM, Prevalent Cases of Obesity, By Age, Both Sexes, N (Millions), Row (%), 2012
- Table 14: 9MM, Prevalent Cases of Obesity, Ages =18 Years, by Sex, N (Millions), Row (%), 2012
- Table 15: 9MM, Prevalent Cases of Obesity Class I, Ages =18 Years, Both Sexes, N (Millions), 2012–2022

- Table 16: 9MM, Prevalent Cases of Obesity Class I, by Age, Both Sexes, N (Millions), Row (%), 2012
- Table 17: 9MM, Prevalent Cases of Obesity Class I, Ages =18 Years, By Sex, N (Millions), Row (%), 2012
- Table 18: 9MM, Prevalent Cases of Obesity Class II, Ages =18 Years, Both Sexes, N (Millions), 2012–2022
- Table 19: 9MM, Prevalent Cases of Obesity Class II, By Age, Both Sexes, N (Millions), Row (%), 2012
- Table 20: 9MM, Prevalent Cases of Obesity Class II, Ages =18 Years, By Sex, N (Millions), Row (%), 2012
- Table 21: 9MM, Prevalent Cases of Obesity Class III, Ages =18 Years, Both Sexes, N (Millions), 2012–2022
- Table 22: 9MM, Prevalent Cases of Obesity Class III, By Age, Both Sexes, N (Millions), Row (%), 2012
- Table 23: 9MM, Prevalent Cases of Obesity Class III, Ages =18 Years, By Sex, N (Millions), Row (%), 2012
- Table 24: 5MM\*, Prevalent Cases of Diagnosed Diabetes among Adults with Overweight/Obesity, Both Sexes, N (Millions), 2012 and 2022
- Table 25: 6MM\*, Prevalent Cases of Diagnosed Hypertension among Adults with Overweight/Obesity, Both Sexes, N (Millions), 2012 and 2022
- Table 26: 4MM, Prevalent Cases of Dyslipidemia\* among Adults with Overweight/Obesity, Both Sexes, N (Millions), 2012 and 2022

## **List Of Figures**

- Figure 1: US, Overweight and Obesity Age-Adjusted Prevalence (%), Ages 20–74 Years, Men, 1960–2010
- Figure 2: US, Overweight and Obesity Age-Adjusted Prevalence (%), Ages 20–74 Years, Women, 1960–2010
- Figure 3: France, Overweight and Obesity Prevalence (%), Ages =15 Years, Men, 1997–2006
- Figure 4: France, Overweight and Obesity Prevalence (%), Ages =15 Years, Women, 1997–2006

- Figure 5: Italy, Overweight and Obesity Prevalence (%), Ages =18, Men, 1983–2010
- Figure 6: Italy, Overweight and Obesity Prevalence (%), Ages =18, Men, 1983–2010
- Figure 7: UK, Overweight and Obesity Prevalence (%), Ages =16 Years, Men, 1993–2011
- Figure 8: UK, Overweight and Obesity Prevalence (%), Ages =16 Years, Women, 1993–2011
- Figure 9: Japan, Obesity Prevalence (%), Ages =20 Years, By Sex, 1976–2006
- Figure 10: Brazil, Overweight and Obesity Prevalence (%), Ages =20 Years, Men, 1975–2003
- Figure 11: Brazil, Overweight and Obesity Prevalence (%), Ages =20 Years, Women, 1975–2003
- Figure 12: Canada, Obesity Prevalence (%), Ages =18 Years, Both Sexes, 1978–2008
- Figure 13: 9MM, Prevalent Cases of Overweight, Ages =18 Years, Both Sexes, N (Millions), 2012–2022
- Figure 14: 9MM, Prevalent Cases of Overweight, by Age, Both Sexes, N (Millions), 2012
- Figure 15: 9MM, Prevalent Cases of Overweight, Ages =18 Years, by Sex, N (Millions), 2012
- Figure 16: 9MM, Overweight Age-Standardized Prevalence (%), Ages =18 Years, By Sex, 2012
- Figure 17: 9MM, Prevalent Cases of Obesity, Ages =18 Years, Both Sexes, N (Millions), 2012–2022
- Figure 18: 9MM, Prevalent Cases of Obesity, By Age, Both Sexes, N (Millions), 2012
- Figure 19: 9MM, Prevalent Cases of Obesity, Ages =18 Years, by Sex, N (Millions), 2012
- Figure 20: 9MM, Obesity Age-Standardized Prevalence (%), Ages =18 Years, By Sex, 2012
- Figure 21: 9MM\*, Prevalent Cases of Obesity Class I, Ages =18 Years, Both Sexes, N (Millions), 2012–2022

- Figure 22: 9MM\*, Prevalent Cases of Obesity Class I, By Age, Both Sexes, N (Millions), 2012
- Figure 23: 9MM\*, Prevalent Cases of Obesity Class I, Ages =18 Years, by Sex, N (Millions), 2012
- Figure 24: 9MM\*, Obesity Class I Age-Standardized Prevalence (%), Ages =18 Years, by Sex, 2012
- Figure 25: 9MM\*, Prevalent Cases of Obesity Class II, Ages =18 Years, Both Sexes, N (Millions), 2012–2022
- Figure 26: 9MM\*, Prevalent Cases of Obesity Class II, By Age, Both Sexes, N (Millions), 2012
- Figure 27: 9MM\*, Prevalent Cases of Obesity Class II, Ages =18 Years, by Sex, N (Millions), 2012
- Figure 28: 9MM\*, Age-Standardized Prevalence (%) of Obesity Class II, Ages =18 Years, By Sex, 2012
- Figure 29: 9MM\*, Prevalent Cases of Obesity Class III, Ages =18 Years, Both Sexes, N (Millions), 2012–2022
- Figure 30: 9MM\*, Prevalent Cases of Obesity Class III, By Age, Both Sexes, N (Millions), 2012
- Figure 31: 9MM\*, Prevalent Cases of Obesity Class III, Ages =18 Years, By Sex, N (Millions), 2012
- Figure 32: 9MM\*, Obesity Class III Age-Standardized Prevalence (%), Ages =18 Years, By Sex, 2012
- Figure 33: 5MM\*, Prevalent Cases of Diagnosed Diabetes among Adults with Overweight/Obesity, Both Sexes, N (Millions), 2012
- Figure 34: 6MM\*, Prevalent Cases of Diagnosed Hypertension among Adults with Overweight/Obesity, Both Sexes, N (Millions), 2012
- Figure 35: 4MM\*, Prevalent Cases of Dyslipidemia\*\* among Adults with Overweight/Obesity, Both Sexes, N (Millions), 2012